

Updated guideline for the management of upper respiratory tract infections in South Africa: 2008

Working Group of the Infectious Diseases Society of Southern Africa
Correspondence to: Dr AJ Brink, email: brinka@ampath.co.za

Introduction: Inappropriate use of antibiotics for non-severe upper respiratory tract infections (URTIs), many of which are viral, adds to the burden of antibiotic resistance. Antibiotic resistance is increasing in *Streptococcus pneumoniae*, the microorganism responsible for most cases of acute otitis media (AOM) and acute bacterial sinusitis (ABS).

Method: The Infectious Diseases Society of Southern Africa held a multidisciplinary meeting to draw up a national guideline for the management of URTIs in 2003. Background information reviewed included randomised controlled trials, existing URTI guidelines and local antibiotic susceptibility patterns. The initial document was drafted at the meeting. Subsequent drafts were circulated to members of the working group for modification. The guideline was published in the South African Medical Journal in 2004 and was a consensus document based upon the opinions of the working group. In 2008 it was decided to update and republish the guideline. This was done electronically using the same working group members, including overseas experts.

Output: Penicillin remains the drug of choice for tonsillopharyngitis. Single-dose parenteral administration of benzathine penicillin is effective, but many favour oral administration twice daily for 10 days. Amoxycillin remains the drug of choice for both AOM and ABS. A dose of 90 mg/kg/day is recommended in general, which should be effective for pneumococci with high-level penicillin resistance (this is particularly likely in children ≤ 2 years of age, in day-care attendees, in cases with prior AOM within the past six months, and in children who have received antibiotics within the last three months). Alternative antibiotic choices are given in the guideline with recommendations for their specific indications. These antibiotics include amoxycillin-clavulanate, some cephalosporins, the macrolide/azalide and ketolide groups of agents and the respiratory fluoroquinolones.

Conclusion: The guideline should assist rational antibiotic prescribing for URTIs. However, it should be continuously updated when new information becomes available from randomised controlled trials and surveillance studies of local antibiotic susceptibility patterns.

Updated and reprinted from S Afr Med J 2004;94(6):475-483 with permission.

SA Fam Pract 2009;51(2):105-114

1. Introduction

There is a worldwide increase in antibiotic resistance, and studies suggest that inappropriate use of antibiotics, particularly for upper respiratory tract infections (URTIs) adds to the burden.¹ Viral infections cause the majority of URTIs. All clinicians should know the natural history of the 'common cold' so that a deviation from normal is managed effectively. Most important is an appreciation that clear nasal secretions frequently become purulent without signifying secondary bacterial disease, and that coughing is a normal accompaniment (Figure 1). The organisms responsible for most bacterial URTIs are similar in all age groups. *Streptococcus pneumoniae* is by far the commonest organism causing otitis media and sinusitis. *Streptococcus pyogenes* is the only significant bacterial

cause of pharyngitis. Systematic reviews suggest that in developed countries the benefit of antibiotics for pharyngitis and otitis media is extremely limited; however, there are few data from poorer countries where rheumatic fever and suppurative complications such as mastoiditis are more common.^{2,3}

South African experience suggests a high prevalence of rheumatic fever in rural areas with poor access to care, an observation in keeping with poor notification rates. For example, 66 new cases of rheumatic heart disease are referred annually to the paediatric cardiology clinic at the Johannesburg Hospital (personal communication — Dr W Hendson). In the Western Cape, 49 new cases are seen every year at Red Cross War Memorial and Tygerberg Children's hospitals (personal communication — Dr J Lawrenson and Prof PL van der




Figure 1: Natural history of the common cold

Merwe). At Umtata General Hospital, 125 children with rheumatic fever were admitted between January 1998 and January 2000, comprising 2% of paediatric admissions. The majority (76%) had two or more episodes of rheumatic fever, which was strongly associated with overcrowding (>six people per room) (personal communication — Proff AS Savio and A Targonski, University of Transkei).

By establishing simple guidelines recommending antibiotics with a relatively narrow spectrum, patients ought to be well managed and serious complications avoided. The most frequently recommended first-line antibiotics for URTIs remain penicillin and amoxicillin. The recommendations for duration of therapy differ; pharyngotonsillitis and acute bacterial sinusitis (ABS) should be treated for 5-10 days and acute otitis media (AOM) for 5-7 days. In this regard, recent evidence suggests that a shorter duration of antibiotic treatment is associated with less emergence of resistant pathogens.⁴ The recommendations for frequency of administration vary according to the site of infection and the pharmacokinetic/pharmacodynamic (PK/PD) profiles of the drugs used; in AOM a twice-daily dose of amoxicillin has the same clinical efficacy as three times a day.

For optimal clinical success, the antibiotic dosage must be tailored to the individual. The most common cause of treatment failure and antibiotic resistance is sub-optimal dosing. For example in AOM, 5 ml is erroneously prescribed as a standard dose for a child weighing 5-15 kg instead of individualising doses by body mass. Dosages given in this guideline include both the registered standard doses and higher doses, which are recommended for use in situations where high-level antibiotic resistance has been reported. Except for amoxicillin and amoxicillin-clavulanate, all paediatric doses are given as mg per kg per dose followed by frequency of daily administration.

Recommendations have been made based on national surveillance of appropriate pathogens and relevant publications.⁵

For *S. pneumoniae*, the most common pathogen causing otitis media and acute sinusitis, resistance to β -lactam antibiotics can be overcome by increasing dosage. For example, a higher dose of amoxicillin of 90 mg/kg/day is generally recommended for treatment of AOM. Because of concerns about the existence of macrolide resistance among isolates of *S. pneumoniae* in some areas of practice in South Africa, in those circumstances this class should preferably be reserved for patients with severe β -lactam antibiotic allergy.

The guideline gives indications for recommended first-line agents as well as alternative choices of antibiotics. The first-line antibiotics are the agents of choice and remain penicillin or amoxicillin. The indications for alternative antibiotics may include the following:

- Allergy or intolerance to first-line agents.
- Recent prior use of first-line agents.
- High-risk cases likely or known to be infected with highly resistant organisms. Consideration needs to be given to β -lactam and macrolide resistance.
- Failed initial therapy.

Few guidelines have been subjected to the rigours of prospective evaluation and most are based on best practice, taking into account unique local circumstances. In South Africa, widespread implementation of throat cultures for pharyngitis is unlikely to occur because an extensive infrastructure would need to be established for an easily treated condition.

2. Acute pharyngotonsillitis

Pharyngotonsillitis is an inflammatory condition of the pharyngeal wall, sometimes divided into pharyngitis and tonsillitis. Respiratory viruses are the major cause of pharyngitis and include the Epstein-Barr virus that causes infectious mononucleosis. Bacteria, especially group A β -haemolytic streptococci (GABHS) (*S. pyogenes*), account for between 5% and 30% of cases. In terms of the ability of throat cultures to diagnose streptococcal pharyngotonsillitis, the specificity approaches 80% while the sensitivity is only 30%, when compared with a rise in or a high level of antistreptolysin O titre.⁶ Non-infectious causes of pharyngitis include allergy and exposure to irritating substances.

Although in most instances GABHS tonsillitis is usually self-limited, antibiotics are recommended to prevent the suppurative and non-suppurative (acute rheumatic fever, glomerulonephritis) post-streptococcal sequelae. The only known risk factor for rheumatic fever is pharyngitis caused by rheumatogenic strains of GABHS, common in both poor rural and urban environments in South Africa. The recommendations of the guideline should therefore be easy to apply and should be such that they would be associated with prevention of rheumatic fever.

2.1 Treatment of streptococcal pharyngotonsillitis

The treatment of choice is penicillin.

2.1.1 Penicillin

There is evidence that intramuscular penicillin prevents rheumatic fever.⁷ Although intramuscular injections are not popular, attempts should be made to administer penicillin parenterally if compliance

and follow-up are unlikely. The mixture of benzathine penicillin and procaine penicillin results in a less painful injection.⁸ When given by mouth, penicillin can be given twice or three times daily instead of four times a day, as previously recommended. Penicillin should be given 30 minutes before a meal as food may reduce its absorption. A 10-day course is still recommended. An important point for continued use of penicillin is the lack of resistance by GABHS, as opposed to erythromycin and other macrolides, where resistance is associated with excessive use.^{9,10} Furthermore, a particular advantage of penicillin is its narrow spectrum of activity.

Children

- Penicillin VK

250 mg twice daily for 10 days (< 27 kg)

500 mg twice daily for 10 days (> 27 kg)

(given 30 minutes before food)

- Benzathine penicillin (intramuscular injection)*

3-5 years: 600,000 U

>5 years: 1.2 MU

Adults and adolescents

- Penicillin VK, 500 mg twice daily for 10 days

(given 30 minutes before food)

- Benzathine penicillin (intramuscular injection),

1.2 MU as single dose*

*Note: To minimise the discomfort of parenteral administration, the medication should be given at room temperature. For patients receiving 1.2 million U, 300,000 U can be given as procaine penicillin.¹¹

2.1.2 Amoxicillin

Amoxicillin is an alternative to penicillin VK and has the advantage of no food restrictions. However, a rash can occur where pharyngotonsillitis is due to Epstein-Barr virus infection. This can lead to an erroneous diagnosis of penicillin allergy or, rarely, a severe skin reaction. Therefore amoxicillin should preferably only be used when GABHS has been identified by culture.

Several recent randomised comparative trials demonstrated non-inferiority of once-daily amoxicillin versus twice-daily amoxicillin or penicillin V.^{11,12} In this regard, an orally administered antibiotic for the treatment of GABHS pharyngitis that could be given once daily would be convenient, particularly for children attending school and may also result in improved compliance.

Children

- <30kg, amoxicillin, 750 mg once daily for 10 days

- >30kg, amoxicillin 1,500mg once daily for 10 days

Adults

- Amoxicillin, 500 mg twice daily for 10 days

2.1.3 Short-course therapy (3-5 days)

Short-course therapy is a recent development.¹³⁻¹⁹ Adam and colleagues¹⁴ showed equivalence between penicillin given three times daily for 10 days and a number of regimens (including use of new macrolides or second-generation cephalosporins) given for five days.¹⁴ The list of agents that have been used for short-course therapy is given below.

Children

- Amoxicillin-clavulanate, 40 mg/kg/day in three divided doses

- Azithromycin, 10-20 mg/kg once daily for three days

- Clarithromycin, 7.5 mg/kg twice daily

- Cefpodoxime, 4 mg/kg twice daily

- Cefprozil 7.5 mg/kg twice daily

- Cefuroxime, 10 mg/kg twice daily

Adults

- Amoxicillin-clavulanate, 375 mg three times daily

- Azithromycin, 500 mg once daily for three days

- Clarithromycin (modified release), 500 mg once daily

- Cefpodoxime, 100 mg twice daily

- Cefprozil, 500mg twice daily

- Cefuroxime, 250 mg twice daily

- Telithromycin, 800 mg once daily

3. Acute otitis media

AOM is one of the most common childhood illnesses — an estimated 75% of children have had more than one episode by 3 years of age. The prevalence is increasing in children less than 2 years of age, and in those attending day-care facilities or exposed to passive smoking.²⁰

The main bacterial causes of AOM are *S. pneumoniae*, non typable *Haemophilus influenzae*, *S. pyogenes* and *Moraxella catarrhalis*. While there is a high rate (~60-70%) of spontaneous resolution of AOM caused by *H. influenzae* and *M. catarrhalis*, *S. pneumoniae* infection is the least likely to resolve spontaneously and therefore the most important target for antibiotic therapy. Antibiotic resistance in *S. pneumoniae* is often clinically relevant because of the relatively poor permeability of antibiotics into the middle ear fluid (MEF).

While most of the drugs approved for AOM have good *in vitro* activity against the common AOM pathogens, there are many differences in their *in vivo* activity and penetration into the MEF. Studies in which efficacy of antibiotics used for AOM is measured by symptomatic relief, fail to discern major differences between drugs because of the so-called 'Pollyanna phenomenon': drugs with poor *in vivo* activity appear to be effective because of the high rate (67%) of spontaneous recovery of the infection.²¹ Given the high rate of spontaneous resolution of AOM, and the probable viral aetiology, particularly in older infants, some authorities argue that antibiotics should be deferred for at least 48 hours during which the patient is given analgesics and decongestants.²² A useful approach may be to dispense the antibiotic or provide a prescription, with the instruction that it is to be given (or the prescription filled) if there has not been resolution by 48 hours. This approach is reasonable where good follow-up is possible in children ≥2 years of age.

However, because of the risks of a serious infection with *S. pneumoniae* and possible limited access of some patients to healthcare in South Africa, we recommend that all cases be treated from the first visit provided that the AOM is correctly diagnosed and distinguished from otitis media with effusion (glue ear); for this reason, the eardrum must be visualised during examination. Because pneumatoscopy is seldom used in South Africa and crying can cause

a red tympanum, AOM is probably seldom correctly diagnosed. In a recent survey of paediatricians attending a training session on otitis media, management from four countries, including South Africa, only half were able to make an accurate otoscopic diagnosis after training.²³ For otitis media associated with a bulging tympanum and a temperature $>38^{\circ}\text{C}$, immediate treatment should be considered. Another reason for immediate antibiotic use is inability to predict whether the patient will have spontaneous resolution.

Antibiotics are essential if AOM is diagnosed in the following patients:

- Recurrent AOM
- Immunocompromised patients
- Neonates
- Structural ENT or immunological abnormalities
- Fever (temperature $>38^{\circ}\text{C}$) or pain >48 hours
- Day-care attendees or siblings of children attending day-care centres

Risk factors for resistant *S. pneumoniae* infections include age (≤ 2 years), attendance at day-care centres or siblings of children attending day-care centres, not vaccinated with the 7-valent pneumococcal conjugate vaccine (PCV), prior AOM within the past six months, and receipt of antibiotics within the last three months. These influence the choice and dosage of antibiotics. Paracetamol (10-15 mg/kg 4-6-hourly) or ibuprofen (10 mg/kg 8-hourly) should be given for analgesia (under- and overdosing are common). Although decongestants are widely prescribed for rhinitis, their use in AOM is controversial. If used, topical application for a maximum of three days is preferable to oral administration.

Otitis media may be part of neonatal sepsis and any neonate with fever should be evaluated for sepsis. Investigations should include a blood culture, urine microscopy, culture and sensitivity testing and lumbar puncture in cases where hypothermia (temperature $<35^{\circ}\text{C}$) or pyrexia (temperature $>38^{\circ}\text{C}$) have been noted. Possible causative organisms in AOM include coliforms, group B streptococci and *Staphylococcus aureus*. Where neonatal sepsis has been excluded and oral therapy is indicated for treatment of AOM, amoxicillin-clavulanate should be given.²⁴ In a recent Israeli study of infants under 2 months of age in whom tympanocentesis had been done, the spectrum of pathogens cultured was similar to that in older children. *S. pneumoniae* was the most common isolate (46%), of which 20% were not susceptible to penicillin. For this reason, the use of a higher dose of amoxicillin (90 mg/kg/day) or amoxicillin-clavulanate, plus additional amoxicillin (to a total dose of amoxicillin of 90 mg/kg/day), would be appropriate.²⁵

Acute otitis media with tympanostomy tubes (AOMT)

AOM with otorrhoea in patients with tympanostomy tubes has not previously been considered as a separate clinical entity in guidelines for the management of upper respiratory tract infections. However, this is a common problem and culture of otorrhoea fluid will often show significant growth of bacteria such as *Pseudomonas aeruginosa* and *S. aureus* in addition to the more usual pathogens, namely *S. pneumoniae*, *H. influenzae* and *M. catarrhalis*. Oral treatment with β -lactams in the presence of the former pathogens would therefore be inappropriate, and in this regard, a topical otological formulation of ciprofloxacin was recently shown to be an efficacious therapeutic

option for the anti-infective management of AOMT, alleviating the need for oral treatment.²⁶ The advantages of topical rather than systemic antibiotics are multiple and include significantly higher tissue levels, substantially reduced adverse effects and perhaps most importantly, considerable less likelihood of the development of antimicrobial resistance.²⁶

3.1 Treatment of AOM

The treatment of choice, except in neonates, is amoxicillin.

3.1.1 Amoxicillin

The dosage recommended is 90 mg/kg/day. This higher dosage is generally recommended since it should provide adequate cover for pneumococcal isolates with high-level penicillin resistance, particularly prevalent in the following circumstances:

- Age ≤ 2 years
- Day-care attendees or siblings of children attending day-care centres
- AOM in previous six months
- Receipt of antibiotics during the three months preceding the AOM episode

At the standard dosage of 40-50 mg/kg/day, amoxicillin is likely to be reasonably effective for an initial episode of AOM in cases infected with non- β -lactamase-producing *H. influenzae* and penicillin-susceptible and possibly intermediate-resistant *S. pneumoniae*. This dosage may not be high enough to eradicate highly penicillin-resistant strains with minimum inhibitory concentrations (MICs) ≥ 2 $\mu\text{g/ml}$.

3.1.2 Amoxicillin-clavulanate

The addition of a β -lactamase inhibitor extends the spectrum of amoxicillin against β -lactamase-producing *H. influenzae* and *M. catarrhalis*. A higher dosage of amoxicillin-clavulanate (90 mg/kg/day of amoxicillin and a constant amount of clavulanate 6.4 mg/kg/day) was previously evaluated in a double-tap study (tympanocentesis at the beginning and end of treatment course). All penicillin-susceptible and intermediate-resistant and 91% of high-level penicillin-resistant pneumococci were eradicated. In addition, 95% of *H. influenzae* and all *M. catarrhalis* were eradicated.²⁷ This formulation is now registered in South Africa and obviates the need to add amoxicillin to amoxicillin-clavulanate to give 90 mg/kg/day of amoxicillin.

3.1.3 Oral cephalosporins

Cefuroxime axetil and cefpodoxime are the only oral cephalosporins that may reach MEF levels sufficiently above the MIC for both penicillin-sensitive and some intermediate-resistant *S. pneumoniae* and for *H. influenzae*.²⁸ Considering the high prevalence of β -lactam resistance in many areas of South Africa, it is recommended that if these cephalosporins are used for treatment of AOM, they should be prescribed at the higher dosages detailed below. The bacterial efficacy of cefpodoxime has not been evaluated in prospective, comparative, double-tap studies. Based upon pharmacokinetic/pharmacodynamic (PK/PD) findings and a clinical trial of otitis media, cefprozil (15 mg/kg twice daily) should not be used empirically in this setting as it is only effective against penicillin-susceptible

pneumococci.^{29,30} Cefaclor, cefixime and loracarbef are less active *in vitro* against *S. pneumoniae* and are not recommended.

3.1.4 Parenteral cephalosporins

The MEF concentration of ceftriaxone exceeds the MICs for AOM pathogens for >50 hours after a single 50 mg/kg intramuscular injection. However, a three-day regimen is clinically superior, particularly in non-responsive AOM caused by penicillin-resistant *S. pneumoniae*.³¹ Ceftriaxone use should be restricted to cases with failure of high-dose amoxicillin-clavulanate or for severe presentations.

3.1.5 Macrolides/azalide/clindamycin

Erythromycin should not be used for the empiric treatment of AOM owing to substantial resistance of *S. pneumoniae* and *H. influenzae* and sub-optimal PK/PD parameters. Azithromycin and clarithromycin have good *in vitro* activity against macrolide-susceptible pneumococci and *M. catarrhalis* and acceptable activity against *H. influenzae*. However, double-tap studies have recently demonstrated that azithromycin at standard doses failed to eradicate macrolide-resistant pneumococci and *H. influenzae*.³² In areas with a high prevalence of macrolide resistance macrolides should therefore preferably be restricted to cases with β -lactam allergy, or used for suspected or proven infections with 'atypical' pathogens. Clindamycin retains activity against most penicillin-resistant *S. pneumoniae* but has no activity against *H. influenzae* or *M. catarrhalis*. It can be used in confirmed, clindamycin-susceptible, pneumococcal AOM, unresponsive to β -lactam antibiotics.

3.1.6 Trimethoprim-sulfamethoxazole (TMP-SMX)

The high rate of resistance of *S. pneumoniae* in South Africa precludes the use of TMP-SMX. High bacteriological failure rates have been noted to occur in double-tap studies.³³

3.2 Duration of therapy for AOM

Most antibiotics are clinically effective for uncomplicated AOM when used in regimens of 5-7 days, since eradication of organisms takes place within 72 hours.³⁴ However, therapy beyond 72 hours is required for adequate eradication of potentially pathogenic bacteria colonising the nasopharynx, because if these are not successfully eradicated they may predispose to relapses of AOM. Further studies are needed to determine optimal duration of therapy in children younger than 2 years of age and in patients with non-responsive AOM.^{28,35} Until then, therapy for 7-10 days is recommended for AOM in the following groups:

- Age ≤ 2 years
- Recurrent or chronic AOM
- Complicated AOM

3.3 Failure to respond to antibiotics

In cases of clinical failure (eg. persistent fever) after 72 hours of appropriate, compliant initial antibiotic therapy, consider referral to an otorhinolaryngologist for tympanocentesis and MEF culture. This is of relevance in areas with a high prevalence of antibiotic-resistant *S. pneumoniae*, as is the case for the majority of major urban centres in South Africa, particularly in the private sector.

3.4 Antibiotic recommendations for AOM

3.4.1 Children

First-line recommended therapy:

- Amoxicillin, 90 mg/kg/day into two or three divided doses for 5-7 days.

Alternative antibiotic choices:

(a) Beta-lactamase-stable antibiotics:

- Amoxicillin-clavulanate, 90 mg/kg/day total dose of amoxicillin divided into two or three doses for 5-7 days
- Cefpodoxime proxetil, 8-16 mg/kg twice daily for 5-7 days
- Cefuroxime axetil, 15-30 mg/kg twice daily for 5-7 days

The higher dosages of cephalosporins recommended would cover for most pneumococcal isolates of intermediate resistance to penicillin, but not necessarily for pneumococcal isolates with high-level resistance. The particular choice of cephalosporins would depend on physician or patient preference, availability and cost. Risk factors for AOM caused by β -lactamase-producing pathogens may include immunocompromised patients and/or neonates.

(b) Antibiotics for β -lactam allergy:

- Azithromycin, 10 mg/kg once daily for three days*
- Clarithromycin, 7.5-15 mg/kg twice daily for 5-7 days*
- Erythromycin estolate, 40 mg/kg twice daily for 5-7 days*
- Cefpodoxime proxetil, 8-16 mg/kg twice daily for 5-7 days†
- Cefuroxime axetil, 15-30 mg/kg twice daily for 5-7 days†

*Macrolides/azalide are recommended for patients who have severe β -lactam allergy.
†Cephalosporins may be considered initially for patients with penicillin intolerance/nontype 1-hypersensitivity reactions (eg. rash).

(c) Failed initial therapy:

- Amoxicillin-clavulanate, with a total dose of 90 mg/kg/day of amoxicillin, divided into two or three doses for 5-7 days for failed initial therapy with amoxicillin alone
- Ceftriaxone, intravenous (IV) or intramuscular (IM), 50-75 mg/kg once daily for three days. This is also recommended in the case of isolates of known high-level antibiotic resistance and in severe presentations, eg. threatened mastoiditis, preferably in consultation with an otorhinolaryngologist.

3.4.2 Adults

The treatment options for AOM in adults are the same as for acute bacterial sinusitis.

3.5 Treatment of AOMT

- Ciprofloxacin 0.3%/dexamethasone 0.1% otic suspension, four drops twice-daily for seven days*

* The technique of application of the drops is essential to the success of this treatment regimen. Aural toilette by suctioning must be performed prior to instilling the drops into the external auditory canal and must be followed by tragal pressure to push the drops through the tympanostomy tubes into the middle ear. A disposable nasal aspirator may be used for suctioning at home.

3.6 Impact of the 7-valent pneumococcal conjugate vaccine (PCV) on otitis media

The most significant development since the original guideline was published has been the introduction of the 7-valent PCV, which has

had a significant impact on the prevalence of otitis media. In this regard, the effectiveness has been monitored in the USA since its deployment in 2001. National rates for AOM visits have decreased by 20% in infants and children <2 years of age, representing 246 fewer AOM visits per 1,000.³⁶ Another national study, comparing 2004 rates with 1997-1999 rates, found that ambulatory visits and antibiotic prescriptions attributable to AOM decreased from 2,173 to 1,244 visits per 1,000 person-years (42.7% reduction) and from 1,244 to 722 prescriptions per 1,000 person-years (42.9% reduction), respectively.³⁷

In Tennessee and New York, comparing the 2000-2001 birth cohort to the 1998-1999 birth cohort, frequent AOM (three episodes in six months or four episodes in one year) in children declined by 17% and 28%, and pressure-equalising tube insertions declined by 16% and 23%, respectively.³⁸ Acute otitis media and treatment failure in AOM were shown to have decreased by 24% in infants and children in New York coincident with the introduction of high dose amoxicillin therapy and the 7-valent PCV.³⁹ The same researchers subsequently also warned that penicillin non-susceptible *S. pneumoniae* may be re-emerging in that population in children with recurrent and difficult-to-treat AOM.⁴⁰

From a pharmaco-economic point of view, the total estimated US national direct medical expenditure for AOM-related ambulatory visits and antibiotic prescriptions for infants and children <2 years of age decreased from an average of US\$ 1.41 billion during 1997 to 1999 to US\$ 0.95 billion in 2004, a 32.3% reduction in expenditure.³⁷

The recent announcement by the South African Department of Health that it will make the 7-valent pneumococcal conjugate vaccine (PCV) available to all South African children has been hailed by all infectious disease and associated healthcare professionals as a significant step in achieving South Africa's millennium goal of reducing child mortality to 20 deaths per 1,000 children by 2015. This revolutionary vaccine, which was made available in the Eastern Cape at the end of 2008 and will be rolled out nationally this year, is crucial, particularly in the light of the impact of invasive pneumococcal disease in a country with a high burden of HIV infection. Added potential benefits of this vaccine, in addition to its potential impact on the prevalence of otitis media, include its 'herd immunity'. In the light of a different 'epidemiological climate' in South Africa, it is hoped that this vaccine will offer the same benefits as published elsewhere.

4. Acute bacterial sinusitis

ABS is most often preceded by a viral URTI. Allergy, trauma, dental infection, or other factors that lead to inflammation of the nose and paranasal sinuses may also predispose individuals to ABS. The most common bacterial isolates from the maxillary sinuses of patients with ABS are similar to those isolated in AOM, namely *S. pneumoniae*, *H. influenzae* and *M. catarrhalis*. As with AOM, *S. pneumoniae* causes most of the serious sequelae and antibiotic therapy must be capable of eradicating it. Other streptococcal species, anaerobic bacteria and *S. aureus* occur in a small percentage of cases. *Chlamydia pneumoniae* and other 'atypical' pathogens should be considered in patients with chronic sinusitis.⁴¹ Fungi have been associated with sinusitis and may be seen in allergic sinusitis and immunocompromised hosts. However, their clinical significance in immunocompetent patients is unclear.

Multiple factors play a role in the antibiotic selection for ABS. *S. pneumoniae* may be associated with serious intracranial and extracranial complications, so it should be adequately covered in initial therapy. Gram-negative cover for *H. influenzae* (and *M. catarrhalis* in children) cannot be ignored. Prior antibiotic use is a major risk factor for infection with antibiotic-resistant strains. Because recent antibiotic exposure increases the risk of carriage and infection with resistant organisms, the choice and dosage of antimicrobial therapy should take into account a history of recent antibiotic use. Other factors to consider are the severity of disease, its rate of progression and varying rates of resistance within South Africa. However, regarding the issue of whether antibiotics are necessary for ABS, recent meta-analyses of antibiotics versus placebo showed marginal benefit.^{42,43} In future, the Federal Drug Administration of the United States of America will be demanding placebo studies for this indication.

4.1 Duration of antibiotic treatment for ABS

The duration of antibiotic treatment for ABS is 10 days. This is based on published studies of clinical trials in which pretreatment and post-treatment sinus aspirates were performed.⁴⁴ However, evidence for moxifloxacin and telithromycin suggests that a shorter course of 5-7 days is clinically and/or bacteriologically equivalent to a 10-day course.^{45,46} Recent studies have also shown that bacteriological eradication occurs within 72 hours with moxifloxacin (400 mg once daily) or with high-dose, short-course levofloxacin (750 mg once daily for five days).^{47,48} This higher dose of levofloxacin improves the PK/PD profile of the agent and in a comparative trial of this dose versus levofloxacin 500mg once daily for 10 days, clinical and microbiological efficacy was found to be similar.⁴⁹ Gemifloxacin has recently been registered in South Africa as a five-day course for ABS.

4.2 Failure to respond to antibiotics

In cases of clinical failure (eg. persistent fever) after 72 hours of appropriate, compliant antibiotic therapy, consider referral to an otorhinolaryngologist for further evaluation. A computed tomography (CT) scan, fibre-optic endoscopy, or sinus aspiration and culture may be necessary. This is of relevance in areas with a high prevalence of antibiotic-resistant *S. pneumoniae*, as is the case for the majority of major urban centres in South Africa, particularly in the private sector.

4.3 Beta-lactam allergy

Clinicians should differentiate an immediate type-1 hypersensitivity reaction from other less dangerous side effects. Urticaria, angio-oedema and bronchospasm are especially dangerous signs and should contraindicate use of antibiotics that have caused these reactions in a patient in the past. Patients with other types of reactions and side effects may tolerate one specific β -lactam but not another. The cross-reactivity for penicillin and first-generation cephalosporins is higher than for second- or third-generation cephalosporins. Consensus opinion suggests that if the previous reaction to penicillin was an itchy maculopapular rash, it would be relatively safe to use cephalosporins.⁵⁰ This approach may be followed if the allergy to penicillin was mild, the indication for cephalosporins is well motivated and skin testing for penicillin is impractical.

RTIS

HOW DOES YOUR ANTIBIOTIC MEASURE UP?

Factive[®]
A gem amongst quinolones

Fast, Active
FACTIVE[®]
gemifloxacin mesylate



Healthcare. We Care.



S4 Factive[®]. Each film-coated tablet contains gemifloxacin mesylate equivalent to 320 mg gemifloxacin. Reg. No.: 35/20.1.1/0058.

Marketed by Aspen Pharmacare for Pharmacare Ltd. Co. Reg. No.: 1898/000252/06.
Building 12, Healthcare Park, Woodlands Drive, Woodmead, Sandton, 2148, Johannesburg.
Tel.: 0112393400. www.aspenpharma.com Medical Information Hotline: 0800 118 088 A9161 04/08

4.4 Antibiotic recommendations for ABS

4.4.1 Children

First-line recommended therapy:

- Amoxicillin, 90 mg/kg/day into three divided doses for 10 days.

Alternative antibiotic choices:

(a) Beta-lactamase-stable antibiotics:

- Amoxicillin-clavulanate, 90 mg/kg/day total dose of amoxicillin divided into two or three doses for 10 days
- Cefpodoxime proxetil, 8-16 mg/kg twice daily for 10 days
- Cefuroxime axetil, 15-30 mg/kg twice daily for 10 days

The higher dosages of cephalosporins recommended would cover for most pneumococcal isolates of intermediate resistance to penicillin but not necessarily for pneumococcal isolates with high-level resistance. The particular choice of cephalosporins would depend on physician or patient preference, availability and cost. Risk factors for ABS caused by β -lactamase-producing pathogens may include immunocompromised patients and/or neonates.

(b) Antibiotics for β -lactam allergy:

- Azithromycin, 10 mg/kg once daily for three days*
- Clarithromycin, 7.5-15 mg/kg twice daily for 10 days*
- Erythromycin estolate, 40 mg/kg twice daily for 10 days*
- Cefpodoxime proxetil, 8-16 mg/kg twice daily for 10 days*
- Cefuroxime axetil, 15-30 mg/kg twice daily for 10 days*

*Macrolides/azalide are recommended for severe β -lactam allergy in children.

(c) Failed initial therapy:

- Amoxicillin-clavulanate, with a total dose of 90 mg/kg/day of amoxicillin divided into two or three doses for 10 days for failed initial therapy with amoxicillin alone
- Ceftriaxone, IV or IM, 50-75 mg/kg once daily for 3-5 days. This is also recommended in the case of isolates of known high-level antibiotic resistance and in severe presentations, eg. peri-orbital inflammation, preferably in consultation with an otorhinolaryngologist

4.4.2 Adults

First-line recommended therapy:

- Amoxicillin, 1 g three times daily for 10 days.

Alternative antibiotic choices:

(a) Beta-lactamase-stable antibiotics:

- Amoxicillin-clavulanate, 1 g twice daily plus additional amoxicillin 500 mg twice daily or 2 g sustained release (SR) twice daily for 10 days
- Cefpodoxime proxetil, 200-400 mg twice daily for 10 days
- Cefuroxime axetil, 500 mg-1 g twice daily for 10 days

The higher dosages of cephalosporins recommended would cover for most pneumococcal isolates of intermediate resistance to penicillin but not necessarily for pneumococcal isolates with high-level resistance. The particular choice of cephalosporins would depend on physician or patient preference, availability and cost. Risk factors for ABS caused by β -lactamase-producing pathogens may include immunocompromised patients, including pregnant patients and diabetics.

(b) Antibiotics for β -lactam allergy:

- Azithromycin, 500 mg once daily for three days†
- Clarithromycin (modified release), 1,000 mg once daily for 10 days†
- Erythromycin, 500 mg four times daily for 10 days†
- Telithromycin, 800 mg once daily for 5-10 days†
- Cefpodoxime proxetil, 200-400 mg twice daily for 10 days*
- Cefuroxime axetil, 500 mg-1 g twice daily for 10 days*
- Gemifloxacin, 320 mg daily for 5-10 days‡
- Levofloxacin, 500 mg twice daily or 750 mg once daily for 5-10 days‡
- Moxifloxacin, 400 mg once daily for 5-10 days‡
- Clindamycin, 450 mg three times daily for 10 days.§

*Cephalosporins may be considered initially for patients with penicillin intolerance/nontype-1 hypersensitivity reactions (eg. rash).

†Macrolides/azalide, fluoroquinolones and ketolides are alternative recommendations for severe β -lactam allergy in adults.

‡Respiratory fluoroquinolones are recommended for patients who have recently not responded to other therapy or are intolerant of β -lactams. Ciprofloxacin provides inadequate cover for *S. pneumoniae* and is not recommended.

§Clindamycin use is restricted to confirmed pneumococcal ABS unresponsive to β -lactam antibiotics or as additional therapy to provide for anaerobic and *S. aureus* cover, despite the lack of clinical evidence at this time of the safety or efficacy of combination therapy for ABS.

(c) Failed initial therapy:

- Amoxicillin-clavulanate, 1g twice daily plus additional amoxicillin 500 mg twice daily or 2 g SR for 10 days for failed initial therapy with amoxicillin alone
- Respiratory fluoroquinolones:
 - Gemifloxacin, 320 mg daily for 5 days-10 days
 - Levofloxacin, 500 mg twice daily or 750 mg once daily for 5-10 days
 - Moxifloxacin, 400 mg once daily for 5-10 days
- Telithromycin, 800 mg once daily for 5-10 days
- Ceftriaxone, IV or IM, 1-2 g once daily for 3-5 days
- Ceftriaxone or the respiratory fluoroquinolones may also be used as first-line therapy in severe initial presentations, eg. peri-orbital oedema, preferably in consultation with an otorhinolaryngologist.

References

- Jacobs MR. World trends in antimicrobial resistance among common respiratory tract pathogens in children. *Pediatr Infect Dis J* 2003; **22**: S109-S119
- Del Mar C. Managing sore throat: a literature review. *Med J Aust* 1992; **156**: 572-575
- Del Mar CB, Glasziou PP, Spinks AB. Antibiotics for sore throat (Cochrane Review). In: *The Cochrane Library*, Issue 4, 2003. Chichester, UK: John Wiley & Sons, 2003
- Pechere JC. Parameters important in short antibiotic courses. *J Int Med Res* 2000; **28**(Suppl 1): 3A-12A
- Huebner RE, Wasas AD, Hockman M, et al. Bacterial aetiology of non-resolving otitis media in South African children. *J Laryngol Otol* 2003; **117**: 169-172
- Graham A, Fahey T. Sore throat: diagnostic and therapeutic dilemmas. *BMJ* 1999; **319**: 173-174
- Arguedas A, Mohs E. Prevention of rheumatic fever in Costa Rica. *J Pediatr* 1992; **121**: 569-572
- Bass JW. A review of the rationale and advantages of various mixtures of benzathine penicillin G. *Pediatrics* 1996; **97**: 960-963
- Dicuozzo G, Fiscarelli E, Gherardi G, et al. Erythromycin-resistant pharyngeal isolates of *Streptococcus pyogenes* recovered in Italy. *Antimicrob Agents Chemother* 2002; **46**: 3987-3990
- Reinert RR, Lutticken R, Bryskier A, Al-Lahham A. Macrolide-resistant *Streptococcus pneumoniae* and *Streptococcus pyogenes* in the pediatric population in Germany during 2000-2001. *Antimicrob Agents Chemother* 2003; **47**: 489-493
- Clegg HW, Ryan AG, Dallas SD, et al. Treatment of streptococcal pharyngitis with once-daily compared with twice-daily amoxicillin. *Pediatric Inf Dis J* 2006; **25**: 761-767
- Lennon DR, Farrell E, Martin DR, Stewart JM. Once-daily amoxicillin versus twice-daily penicillin V in group A β -haemolytic streptococcal pharyngitis. *Arch Dis Child* 2008; **93**: 474-478
- American Academy of Pediatrics. Group A streptococcal infections. In: Pickering LK, ed. *Red Book: 2003 Report of the Committee of Infectious Diseases*. Elk Grove Village, Ill.: American Academy of Pediatrics, 2003: 578-580
- Adam D, Scholz H, Helmerking M. Short course antibiotic treatment of 4782 culture-proven cases of group A streptococcal tonsillo-pharyngitis and incidence of poststreptococcal sequelae. *J Infect Dis* 2000; **182**: 509-516

15. Pichichero ME, Margolis PA. A comparison of cephalosporins and penicillins in the treatment of group A beta-hemolytic streptococcal pharyngitis: a meta-analysis supporting the concept of microbial copathogenicity. *Pediatr Infect Dis J* 1991; **10**: 275-281
16. Hebblethwaite EM, Brown GW, Cox DM. A comparison of the efficacy and safety of cefuroxime axetil and augmentin in the treatment of upper respiratory tract infections. *Drugs Exp Clin Res* 1987; **13**: 91-94
17. Portier H, Chavanet P, Gouyon JB, et al. Five day treatment of pharyngotonsillitis with cefpodoxime proxetil. *J Antimicrob Chemother* 1990; **26**: 79-85
18. Mehra S, Van Moerkerke M, Welck J, et al. Short course therapy with cefuroxime axetil for group A streptococcal tonsillopharyngitis in children. *Pediatr Infect Dis J* 1998; **17**: 452-457
19. Pichichero ME, Cohen R. Shortened course of antibiotic therapy for acute otitis media, sinusitis and tonsillopharyngitis. *Pediatr Infect Dis J* 1997; **16**: 680-695
20. Hoppe HL, Johnson CE. Otitis media: focus on antimicrobial resistance and new treatment options. *Am J Health Syst Pharm* 1998; **55**(18): 1881-1897
21. Marchant CD, Carlin SA, Johnson CE, et al. Measuring the comparative efficacy of antibacterials for acute otitis media: the "Pollyanna phenomenon". *J Pediatr* 1992; **120**: 72-77
22. Del Mar C, Glasziou P, Hayem M. Are antibiotics indicated as initial treatment for children with acute otitis media? A meta-analysis. *BMJ* 1997; **314**: 1526-1529
23. Pichichero ME. Diagnostic accuracy of otitis media and tympanocentesis skills assessment among paediatricians. *Eur J Clin Microbiol Infect Dis* 2003; **22**: 519-524
24. Bradley JS, Nelson JD. *Nelson's Pocketbook of Pediatric Antimicrobial Therapy*. Lippincott Williams and Wilkins, 2002. www.skyscape.com (accessed March 2004)
25. Turner D, Leibovitz E, Aran A, et al. Acute otitis media in infants younger than two months of age: microbiology, clinical presentation and therapeutic approach. *Pediatr Infect Dis J* 2002; **21**: 669-674
26. Dohar J, Giles W, Roland P, Bikhazi N, Carroll S, Moe R, et al. Topical ciprofloxacin/dexamethasone superior to oral amoxicillin/clavulanic acid in acute otitis media with otorrhea through tympanostomy tubes. *Pediatrics* 2006; **118**: e561-e569
27. Dagan R, Hoberman A, Johnson C, et al. Bacteriologic and clinical efficacy of high dose amoxicillin/clavulanate in children with acute otitis media. *Pediatr Infect Dis J* 2001; **20**: 829-837
28. Craig WA, Andes D. Pharmacokinetics and pharmacodynamics of antibiotics in otitis media. *Pediatr Infect Dis J* 1996; **15**: 255-259
29. Nicolau DP, Sutherland CA, Arguedas A, Dagan R, Pichichero ME. Pharmacokinetics of cefprozil in plasma and middle ear fluid in children undergoing treatment for acute otitis media. *Pediatr Drugs* 2007; **9**: 119-123
30. Pichichero ME, Dagan R, Arguedas A, et al. A multi-center, open-labeled study of cefprozil in children with persistent and recurrent acute otitis media. Abstract G-819. 45th Interscience Conference on Antimicrobial Agents and Chemotherapy, Washington, D. C., December, 2005
31. Leibovitz E, Piglansky L, Raiz S, et al. Bacteriologic efficacy of a three-day intramuscular ceftriaxone regimen in nonresponsive acute otitis media. *Pediatr Infect Dis J* 1998; **17**: 1126-1131
32. Dagan R, Leibovitz E, Fliss DM, et al. Bacteriologic efficacies of oral azithromycin and oral cefaclor in treatment of acute otitis media in infants and young children. *Antimicrob Agents Chemother* 2000; **44**: 43-50
33. Leiberman A, Leibovitz E, Piglansky L, et al. Bacteriologic and clinical efficacy of trimethoprim-sulfamethoxazole for treatment of acute otitis media. *Pediatr Infect Dis J* 2001; **20**: 260-264
34. Ingvarsson L, Lundgren K. Penicillin treatment of acute otitis media in children. A study of the duration of treatment. *Acta Otolaryngol* 1982; **94**: 283-287
35. Leibovitz E, Dagan R. Otitis media therapy and drug resistance. *Infect Med* 2001; **18**: 263-270
36. Grijalva CG, Poehling KA, Nuorti JP, et al. National impact of universal childhood immunization with pneumococcal conjugate vaccine on outpatient medical care visits in the United States. *Pediatrics* 2006; **118**: 865-873
37. Zhou F, Shefer A, Kong Y, Nuorti JP. Trends in acute otitis media-related health care utilization privately insured young children in the United States, 1997-2004. *Pediatrics* 2008; **121**: 253-260
38. Poehling KA, Szilagyi PG, Grijalva CG, et al. Reduction of frequent otitis media and pressure-equalizing tube insertions in children after introduction of pneumococcal conjugate vaccine. *Pediatrics* 2007; **119**: 707-715
39. Casey JR, Pichichero ME. Changes in frequency and pathogens causing acute otitis media in 1995-2003. *Pediatr Infect Dis J* 2004; **23**: 824-828
40. Pichichero ME, Casey JR, Hoberman A, Schwartz R. Pathogens causing recurrent and difficult-to-treat acute otitis media, 2003-2006. *Clin Pediatr* June 2008 (ePub ahead of print)
41. Dunbar LM. Current issues in the management of bacterial respiratory tract disease: the challenge of antibacterial resistance. *Am J Med Sci* 2003; **326**: 360-368
42. Young J, De Sutter A, Merenstein D, et al. Antibiotics for adults with clinically diagnosed acute rhinosinusitis: a meta-analysis of individual patient data. *Lancet* 2008; **371**: 908-914
43. Ioannidis JPA, Chew P, Lau J. Standardized retrieval of side effects data for meta-analysis of safety outcomes. A feasibility study in acute sinusitis. *J Clin Epidemiol* 2002; **55**: 619-626
44. Gwaltney JM Jr, Scheld WM, Sande MA, et al. The microbial etiology and antimicrobial therapy of adults with acute community-acquired sinusitis: a fifteen-year experience at the University of Virginia and review of other selected studies. *J Allergy Clin Immunol* 1992; **90**: 457-461
45. Siegert R, Gehanno P, Nikolaidis P, et al. A comparison of the safety and efficacy of moxifloxacin (BAY 12-8039) and cefuroxime axetil in the treatment of acute bacterial sinusitis in adults. The Sinusitis Study Group. *Respir Med* 2000; **94**: 337-344
46. Roos K, Brunswig-Pitschner C, Kostrica R, et al. Efficacy and tolerability of once-daily therapy with telithromycin for 5 or 10 days for the treatment of acute maxillary sinusitis. *Chemotherapy* 2002; **48**: 100-108
47. Ariza H, Rojas R, Johnson P, et al. Eradication of common pathogens at days 2, 3 and 4 of moxifloxacin therapy in patients with acute bacterial sinusitis. *BMC Ear, Nose and Throat Disorders* 2006; **6**: 8-15
48. Anon JB, Paglia M, Xiang J, Ambrose PG, Jones RN, Kahn JB. Serial sinus aspirate samples during high-dose, short-course levofloxacin treatment of acute maxillary sinusitis. *Diagn Microbiol Inf Dis* 2007; **57**: 105-107
49. Poole M, Anon J, Paglia M, Xiang J, Khashab M, Kahn J. A trial of high-dose, short-course levofloxacin for the treatment of acute bacterial sinusitis. *Otolaryngology- Head and Neck Surgery* 2006; **134**: 10-17
50. Kelkar PS, Li JT. Cephalosporin allergy. *N Engl J Med* 2001; **345**: 804-809

Working Group of the Infectious Diseases Society of Southern Africa

AJ Brink (Ampath National Laboratory Services, Johannesburg), MF Cotton (Department of Paediatrics and Child Health, Faculty of Health Sciences, University of Stellenbosch and Tygerberg Children's Hospital, Cape Town), C Feldman (Division of Pulmonology, Department of Medicine, Johannesburg Hospital and University of the Witwatersrand, Johannesburg), H Finlayson (Department of Paediatrics and Child Health, Faculty of Health Sciences, University of Stellenbosch and Tygerberg Children's Hospital, Cape Town), L Geffen (SA Academy of Family Practice/Primary Care, Cape Town), R Green (Department of Paediatrics and Child Health, University of Pretoria), W Hendson (Paediatric Cardiology, Department of Paediatrics and Child Health, University of the Witwatersrand, Johannesburg), MH Hockman (Otorhinolaryngologist, Linksfield Park Clinic, Johannesburg), G Maartens (Division of Infectious Diseases, Department of Medicine, University of Cape Town), SA Madhi (NICD/MRC/Wits Respiratory and Meningeal Pathogens Research Unit and Paediatric Infectious Diseases Research Unit, Johannesburg) M Mutua-Mpungu (Department of Family Medicine, MEDUNSA, Pretoria), GH Swingler (Faculty of Health Sciences, School of Child and Adolescent Health, University of Cape Town and Red Cross Children's Hospital, Cape Town).

International review panel

Keith P Klugman (Professor of Infectious Diseases, Department of International Health, The Rollins School of Public Health, Emory University, Atlanta, USA), Ron Dagan (Professor of Paediatric Infectious Diseases, Soroka University Medical Center, Beer-Sheva, Israel), Adriano Arguedas (Professor of Pediatrics, Instituto de Atencion Pediatrica and Universidad de Ciencias Medicas, San Jose, Costa Rica).

Disclosure statement

| Author | Entity, Relationship code |
|----------------|--|
| AJ Brink | Abbott SB AB RC RS; Aventis SB RG RS; Bayer SB; Bristol-Myers Squibb SB RG RC; Merck SB AB RC; Pfizer SB; Roche SB |
| MF Cotton | Abbott SB; Aventis SB; Bristol-Myers Squibb SB GI; Roche SB |
| C Feldman | Abbott AB SB; Astra-Zeneca ES; Aventis AB SB ES RR; Bayer SB; Bristol-Myers Squibb AB ES SB; Merck AB ES; Pfizer ES SB; Roche SB; GlaxoSmithKline ES SB |
| H Finlayson | None |
| L Geffen | None |
| R Green | AstraZeneca AB ES SB; GlaxoSmithKline ES SB; MSD AB ES SB; Pfizer SB, Roche AB SB; Schering-Plough AB SB |
| W Hendson | None |
| MH Hockman | Aventis C AB SB ES; Bayer ES AB SB; Bristol-Myers Squibb AB SB; Pfizer SB; Roche ES SB; GlaxoSmithKline ES SB RS; Servier ES C SB; Schering-Plough ES SB |
| G Maartens | Abbott SB, Aventis SB, GlaxoSmithKline SB, Merck SB, Pfizer SB, Roche SB |
| S A Madhi | Abbott SB RC; Aventis SA; GlaxoSmithKline SB; Wyeth RC RG |
| M Mutua-Mpungu | None |
| G H Swingler | None |

Relationship code: AB = Advisory Board; B = Board member; CL = Collaborator; C = Consultant; EG = Educational grant; ES = Educational support; GI = Grant investigator; I = Investigator; LA = Licensing agreement; RC = Research contractor; RG = Research grant; RR = Research relationship; RS = Research support; SA = Scientific advisor; S = Shareholder; SB = Speakers bureau; U = Unknown.

Summary: Antibiotic recommendations for upper respiratory tract infections

| SORE THROAT | ACUTE OTITIS MEDIA (AOM) | ACUTE BACTERIAL SINUSITIS (ABS) |
|---|--|--|
| Clinical diagnosis of streptococcal pharyngotonsillitis | Clinical diagnosis of AOM | Clinical diagnosis of ABS |
| <p>Points in favour of empiric antimicrobial treatment:</p> <ul style="list-style-type: none"> Acute onset Temperature >38°C Tender anterior cervical nodes Tonsillar erythema/exudates Age 3-15 years Previous rheumatic fever or rheumatic heart disease <p>Points against empiric antimicrobial treatment:</p> <ul style="list-style-type: none"> Rhinorrhoea Cough Diarrhoea Conjunctivitis Age >45 years | <ul style="list-style-type: none"> Visualisation of the eardrum is essential for diagnosis Bulging, red or yellow tympanic membrane Ancillary features include: <ul style="list-style-type: none"> Otalgia Temperature >38°C <p>(Note: an effusion alone is not an indication for antimicrobials)</p> | <p>Consider in adults or children with an upper respiratory tract infection that is NOT improving after 10 days or worsens after 5-7 days and is accompanied by some or all of these symptoms:</p> <ul style="list-style-type: none"> Fever Facial tenderness, particularly unilateral or focused in the region of a sinus group (peri-orbital, maxillary, frontal or ethmoidal) Dental tenderness Nasal discharge, nasal congestion, anosmia (loss of sense of smell), cough, ear fullness and pressure (frontal sinusitis does not occur in toddlers <4 years because of delayed development of the frontal sinuses) |
| Treatment of choice | Treatment of choice | Treatment of choice |
| <p>The treatment of choice is penicillin:</p> <p><i>Children:</i></p> <ul style="list-style-type: none"> Penicillin VK <ul style="list-style-type: none"> 250 mg twice daily for 10 days (≤27 kg) 500 mg twice daily for 10 days (>27 kg) Benzathine penicillin (intramuscular injection)* <ul style="list-style-type: none"> - 3-5 yrs: 600,000 U - >5 yrs: 1.2 MU <p><i>Adults and adolescents:</i></p> <ul style="list-style-type: none"> Penicillin VK, 500 mg twice daily for 10 days Benzathine penicillin (intramuscular injection)* 1.2 MU <p>For patients with severe β-lactam allergy:</p> <p><i>Children:</i></p> <ul style="list-style-type: none"> Erythromycin estolate, 40 mg/kg twice daily for 10 days Azithromycin, 10-20 mg/kg once daily for three days Clarithromycin, 7.5-15 mg/kg twice daily for five days <p><i>Adults:</i></p> <ul style="list-style-type: none"> Erythromycin estolate, 500 mg twice daily for 10 days Azithromycin, 500 mg once daily for three days Clarithromycin (modified release), 500 mg once daily for five days Telithromycin, 800 mg once daily for five days | <p>The treatment of choice is oral amoxicillin:</p> <p><i>Children†:</i></p> <ul style="list-style-type: none"> Amoxicillin, 90 mg/kg/day in two or three divided doses <p><i>Adults:</i></p> <ul style="list-style-type: none"> Amoxicillin, 1,000 mg three times daily for five days <p>For patients with severe β-lactam allergy:</p> <p><i>Children:</i></p> <ul style="list-style-type: none"> Erythromycin estolate, 40 mg/kg twice daily for 5-7 days Azithromycin, 10 mg/kg once daily for three days Clarithromycin, 7.5-15 mg/kg twice daily for 5-7 days <p><i>Adults:</i></p> <ul style="list-style-type: none"> Erythromycin estolate, 500 mg qid for 5-7 days Azithromycin, 500 mg once daily for three days Clarithromycin (modified release), 500 mg or 1 g once daily for 5-7 days | <p>The treatment of choice is oral amoxicillin:</p> <p><i>Children:</i></p> <ul style="list-style-type: none"> Amoxicillin, 90 mg/kg/day in three divided doses for 10 days <p><i>Adults:</i></p> <ul style="list-style-type: none"> Amoxicillin, 1 g three times daily for 10 days <p>For patients with severe β-lactam allergy:</p> <p><i>Children:</i></p> <ul style="list-style-type: none"> Erythromycin estolate, 40 mg/kg twice daily for 10 days Azithromycin, 10 mg/kg once daily for three days Clarithromycin, 15 mg/kg twice daily for 10 days <p><i>Adults:</i></p> <ul style="list-style-type: none"> Macrolides: <ul style="list-style-type: none"> Erythromycin, 500 mg four times daily for 10 days Clarithromycin (modified release), 1,000 mg once daily for 10 days Azithromycin, 500 mg once daily for three days Telithromycin, 800 mg once daily for 5-10 days Fluoroquinolones: <ul style="list-style-type: none"> Gemifloxacin, 320 mg daily for 5-10 days Levofloxacin, 500 mg twice daily or 750 mg once daily for 5-10 days Moxifloxacin, 400 mg once daily for 5-10 days Clindamycin, 450 mg three times daily for 10 days |
| Indications for referral | Indications for referral | Indications for referral |
| <p>Local complications:</p> <ul style="list-style-type: none"> Peritonsillar sepsis including quinsy abscess-cellulitis-trismus (asymmetrical peritonsillar swelling) Recurrent infections (four or more episodes/year) Non-response to initial therapy <p>Systemic complications:</p> <ul style="list-style-type: none"> Acute rheumatic fever Severe systemic illness | <ul style="list-style-type: none"> If tympanic membrane is not visualised Non-responsive AOM (no improvement after three days of treatment) Suspected intracranial extension Lower motor neuron VIIth nerve palsy Suspected mastoiditis | <ul style="list-style-type: none"> Failure to respond after 72 hours Peri-orbital swelling Evidence of CNS extension (meningism, focal neurological signs, altered level of consciousness) Severe systemic illness Chronic sinusitis – symptomatic >30 days |
| Special investigations | Special investigations | Special investigations |
| <p>Throat swabs should be reserved for patients with recurrent sore throats</p> | <p>None recommended</p> | <ul style="list-style-type: none"> X-rays are of limited value CT scans should be reserved for cases in which surgery is considered Nasal swabs for microbiological investigation are of no value, and specimens for culture and microscopy should only be collected from sinus puncture <p>*Note: To minimise the discomfort of parenteral administration, the medication should be given at room temperature. For patients receiving 1.2 million U, 300,000 U can be given as procaine penicillin. †Duration of therapy: 5-7 days except ≤2 years of age and complicated cases (7-10 days)</p> |