

Is unemployment a major reason for firearm attacks in rural South Africa?

Ogunbanjo, GA, MBBS, MFGP (SA), M Fam Med (Medunsa), FACRRM
Dept. of Family Medicine & Primary Health Care, Medical University
of Southern Africa (MEDUNSA), Pretoria, South Africa

Sotade, BO, MBBS, M Fam Med (Medunsa)
Formerly of the Church of Scotland Hospital,
Tugela Ferry, KwaZulu-Natal, South Africa

Correspondence to:

Prof. Gboyega A Ogunbanjo
Dept of Family Medicine & Primary Health Care, Medical University of Southern Africa
PO Box. 222, Medunsa 0204 South Africa
E-mail: gboyega@intekom.co.za

(SA Fam Pract 2003;45(10): 18-19)

Keywords: unemployment, firearm attacks, victims, South Africa

Introduction

In South Africa, firearms are increasingly used in interpersonal and factional violence. In a five year period (1987 – 1992), gunshot wounds of the torso increased by 300% in KwaZulu-Natal.¹ During the same period, King Edward VIII Hospital in Durban recorded a mortality rate for firearm-related injuries of eight times that for stab wounds and “direct admissions” to the mortuary, three times as common in cases of gunshot wounds compared with stab wounds.² In our descriptive study, all cases of firearm attacks seen at the Church of Scotland Hospital and the government mortuary at Tugela Ferry in KwaZulu-Natal between December 1998 and May 1999 were reviewed to find out the reasons for the attacks. All patients treated at the hospital for non-fatal firearm injuries were interviewed using a pre-tested questionnaire and records from the district surgeon and police were examined to identify all fatal firearm injuries.

Profile of the victims

One hundred and fifty cases of firearm injuries were identified and reviewed. Seventy-four of these people (49,4%)

sustained non-fatal injuries and were all treated at the hospital, while 76 (50,6%) died from their injuries and were taken to the government mortuary. There were 124 male (82,7%) and 26 female (17,3%) victims. Slightly more than half of the victims (79; 52,7%) were between the ages of 20 and 39, and 95 (63,5%) were unemployed. In addition, there was a preponderance of males among the victims with fatal firearm injuries (68 out of 76; 89,5%).

Reasons for the firearm injuries

The majority of the victims (79,4%) could provide no reasons for the firearm attacks, while only a fifth reported reasons such as car hijack, robbery, interpersonal conflict, taxi violence and attempted suicide (Table I).

The most common causes of death among the fatally injured victims (n = 76) were massive hemorrhage from firearm injury site(s) and brain damage. The authors noted that firearm injuries in this rural setting were predominantly a male phenomenon, with a male to female ratio of 4:1. As a public health issue, the impact of these attacks among males is very striking. The male preponderance might be attributed to a high level of unemployment and increased aggression among males, who are more likely to settle interpersonal conflicts by physical means or by using firearms.^{3,4,5,6}

Undeterminable, intentional assault and homicide have been shown to be the most frequent reasons associated with firearm attacks in urban areas.⁷ In our

Table I: Reasons for firearm injuries

Reasons	Frequency	Percentage
Unknown	119	79,4%
Car hijack	9	6,0%
Robbery	4	2,7%
Interpersonal conflict	14	9,3%
Taxi violence	2	1,3%
Attempted suicide	2	1,3%
Total	150	100%

study, the same could not be said of this rural area, as the majority of victims had no record of the reasons for their attacks. We suspect that the victims with non-fatal injuries were afraid of possible retaliation or reprisal by their assailants and did not report the reasons to the police or doctor when they were seen to. Powell and Tanz noted that families in urban communities with income below the poverty level accounted for a high percentage of assaults.⁸ It seems that our finding of high unemployment among the victims of firearm attacks might be unique to this rural area of South Africa. In this study, firearm attacks seen at Tugela Ferry were predominant amongst unemployed young men between the ages of 20 and 39. It would be important

for future studies to focus on the relationship between unemployment and firearm attacks in the country. We believe that the provision of employment opportunities will help to reduce the incidence of firearm injuries in this rural area, and possibly also in other parts of the country. This is an example of a social issue that has a direct effect on the public health system.

References

1. Muckart DJJ, Meumann C, Botha JBC. The changing pattern of penetrating torso trauma in KwaZulu-Natal – a clinical and pathological review. *SAMJ* 1995;85(11):1172-4.
2. Muckart DJJ. Trauma – the malignant epidemic. *SAMJ* 1991;79:93-5.
3. Azmak D, Altun G, Bilgi S, Yilmaz A. Firearm fatalities in Edime, 1984–1997. *Forensic Sci Int* 1998;95(3):231-9.
4. Elfawal MA, Awad OA. Firearm fatalities in Eastern Saudi Arabia: impact of culture and legislation. *Am J Forensic Med Pathol* 1997;18(4):391-6.
5. Strong RW. Gunshot wounds of Adolescents. *Med J Aust* 1980;1:113-5.
6. Byarugaba J, Kielkowski D. Reflections on trauma and violence-related deaths in Soweto, July 1990-June 1991. *SAMJ* 1994;84(9):610-4.
7. Wigton A. Firearm-related injuries and deaths among children and adolescents in Cape Town 1992–1996. *SAMJ* 1999;89:407-10.
8. Powell EC, Tanz RR. Child and adolescent injury and death from urban firearm assaults: association with age, race, and poverty. *Inj Prev* 1999;5(1):41-7.

MASTERS DEGREE IN FAMILY MEDICINE & PRIMARY HEALTH CARE, MEDUNSA M Med (Family Medicine)

MEDICAL UNIVERSITY
OF SOUTHERN AFRICA



MEDUNSA



The Department of Family Medicine & Primary Health Care Medunsa invites applications from doctors who wish to enroll for its M Med (Family Medicine) program in the year 2004, and on completion register as Family Physicians. The program spans a minimum of four years with regular contact sessions up to five times per year at the department and in provincial groups with local facilitators.

Requirements:

1. Possession of MBChB degree or equivalent qualification for at least two years
2. Clinical work should be done in Primary Health Care settings or district hospitals with a strong association with PHC patients.
3. Registration as a medical practitioner with the Health Professions Council of South Africa or with the regulatory body of the country in which the doctor is practising.

Contact Sessions:

February Contact Week: 2nd – 6th February 2004. Students will be expected to attend contact sessions at Medunsa and in the provincial groups throughout the years. The program covers the following modules: The consultation; Human development (including whole person medicine); The Family; Ethics including screening; Research Methods; Applied Social Science in Medicine; Learning; Portfolio development; Practice management & organisation of health services; Evidence Based Medicine; Clinical reasoning & Therapeutics; Prevention and health promotion; Quality improvement; Principles & Foundations of Family Medicine

Applications close Mid January 2004

Application forms and further information are obtainable from: HOD, Department of Family Medicine & Primary Health Care, PO Box 222, Medunsa 0204

Tel: (012) 521 4314/4528 Fax: (012) 521 4172

e-mail: ansie@wn.apc.org

We empower the educationally disadvantaged community of Southern Africa