CPD Accreditation Numbers:

CPD@SA Family Practice

Earn 100% of your required CEU points by completing the SA FAMILY PRACTICE CPD questionnaire each month.

2006 ANSWERSHEETS

CPD QUESTIONNAIRES MUST NOW BE COMPLETED ONLINE AT www.safpj.co.za

- Simply click on the "CPD" icon below the journal masthead. You will be directed to the

- Simply click on the CPD Icon below the journal mastnead. You will be directed to the CPD questionnaire on the www.ecpdsa.co.za website.

 Click on the SA Family Practice Volume 48(8) September 2006 activity listed on the "Choose a CPD activity" page.

 You will be prompted to login on the ecpdsa website. Click on "Register" on the top menu bar if you have not registered on ecpdsa before, or simply Log In.

 Click on "Continue" at the bottom of the "CPD Activity" page to view the CPD questions. Complete the questions online. If successful, you will receive your CPD actificate police.
- If you experience any problems or choose to receive a printed CPD answer sheet for manual processing, kindly contact Léhane at Tel: (021) 975 4684, Fax: 086 500 4974, or Email: lehane@ecpdsa.co.za If you return the completed CPD answer sheet by fax or email your answers will be entered on the www.ecpdsa.co.za website. If successful, your CPD certificate will be
- available on ecpdsa, or returned to you on request.

Please note this is a FREE service to members of the SA Academy of Family Practice/Primary Care and subscribers to the journal only.

Thank you very much for your cooperation. The CPD Management Office

Question 1-5: Oral Manifestations of Viral Infections: Page 20-24

The infection that accompanies primary herpetic gingivostomatitis is usually subclinical in early adulthood.

- True
- В. False

2. Choose the incorrect statement

- Around 45% of the community is affected by episodes of herpes labialis.
- B. Emotional upset can predispose the patient to recurrent herpes infection
- The lesions of secondary herpes labialis ulcerate and become crusted before heal-C. ing within 10 days.
- There is no drug available to prevent the migration of the herpes virus to the tri-D. geminal ganglion after primary infection.
- Antiviral drugs, such as acyclovir, are only useful if applied during the prodromal stage.

Dental manipulation will not lead to reoccurrence of varicella - zoster virus.

- True
- В.

True or False

The cutaneous vesicles of hand, foot and mouth disease can resemble chickenpox.

- True

True or false?

The gingival and soft palate are the most common locations for squamous cell papilloma.

- В.

Question 6-10: Aboviral Diseases in Southern Africa: Page 25-28

Choose the incorrect statement

- Mosquitoes and midges are arthropods that can transmit viruses to humans.
- Phlebo viruses are generally transmitted by ticks.
- C Mild aseptic meningitis and encephalitis are acute central nervous system illnesses seen with arboviral infections.
- Rift Valley Fever is caused by the bunyaviridae.
- The flaviviridae cause yellow fever.

Rift Valley Fever is the most important viral haemorrhagic fever in South Africa.

- В. False

True or false?

The Aedes aegyptis mosquito transmits the Fogaviridae.

- R

9. True or false?

Sindbis skin rash has a typical "fried egg" appearance.

- В False

Choose the incorrect statement 10.

- Yellow fever is not an endemic disease in South Africa.
- A valid certificate of immunization against yellow fever is valid for 6 years.
- The West Nile virus is widespread in Europe and India.
- Bacterial sepsis is on the list of differential diagnosis for haemorrhagic fever.
- Specific antiviral agents against the arboviridae do not exist.

Question 11-15: Viral Hepatitis: Page 29-34

11. True or false?

Acute viral hepatitis may be asymptomatic.

- True
- False

True or false?

Raw vegetables are a particularly common source of hepatitis A infec-

- В False

True or false?

Person-to-person transmission of hepatitis E is common.

- True
- В.

Choose the incorrect statement

- The hepatitis B virus is an important cause of hepatocellular carcinoma.
- In hepatitis B infection, 70% of patients will remain anicteric.
- Serum aminotransferases may be normal in patients with hepatitis B.
- Patient who have acquired natural immunity to hepatitis B through infection develop both anti-HBs and anti-HBc antibodies.
- Ε Patients who are HBV-DNA negative and HBsAg positive have occult hepatitis

Choose the incorrect statement 15.

- Hepatitis C is one of the main indications for liver transplantation in developed
- Hepatitis C is shed in breast milk, but breastfeeding is not contraindicated.
- The incubation period of hepatitis C varies from 6 to 12 months.
- Cryoglobinaemia, caused by hepatitis C, results in immune complex-mediated vasculitis.
- False positive HCV antibodies may occur in a population with a low prevalence

Question 16-20: The Basics of Viral Vaccines: Page 36-43

Choose the incorrect statement

- Simple saline may be the suspending fluid in vaccines.
- With passive immunisation, preformed antibodies are administered.
- Herd immunity occurs when both active and passive immunization are adminis-
- Thimerosal is a preservative found in vaccines.
- Adjuvants are substances that are added to vaccines to enhance their immunogenicity.

17. True or false?

- Immunoglobulins have no effect on live attenuated virus vaccines.
- В False

18. True or false?

The measles vaccine is administered at 9 and 18 months.

True or false?

Children on cytotoxic therapy should not receive live vaccines.

- True
- В.

20. Choose the incorrect statement

- Inactivated vaccines are not as immunogenic as live attenuated vaccines.
- В. Risk factors for non-response to hepatitis B vaccines include age greater than 15 years and low weight
- Children on long-term aspirin therapy constitute an indication for vaccination against influenza.
- Sewerage workers are at risk for developing hepatitis A infection.
- Accidental administration of rubella vaccine is not regarded as a reason for termination of pregnancy.